

Comments on Dragging Effects

Dieter R. Brill

Department of Physics
University of Maryland
College Park MD 20742 U.S.A.

Submitted to volume in *Einstein Studies* series

Abstract. This is a reply, given at the conference “Mach’s Principle” in Tübingen in July 1993, to the paper by Pfister (1993).

1. Preliminaries and Frivolities

The “Machian” effects of rotation on inertial frames have a distinguished history, as pointed out in the previous paper by Pfister (1993). But beyond that, the appeal of Mach’s ideas has also been used to motivate points of view that often are well-founded in inverse relation to their attraction for the general public. For example, in a recent “popular” scientific book (Fahr 1993), Mach occurs as early as page 55, and there is a whole chapter devoted to matters Machian, with the somewhat ominous title* *The view in the large — doomed to failure?* Even if the answers of this particular book may leave much to be desired, it reminds us that these questions are not esoteric considerations of a select few, but of direct and immediate interest to a large number of intellectuals; and one can only wish that the present volume may have some fraction of the impact that such “popular” books have on the general public!

Pfister’s emphasis is not so much on Mach as an abstract principle, but on definite and calculable effects suggested by the Machian line of thought. This is a very useful and practical approach; it is likely that Mach’s ideas have brought more progress by suggesting calculations of such effects than by any general formulation of his principle. This view has been expressed by Prof. Shimony (1992), who compares Mach to a Yankee storekeeper with many useful items on his shelves, connected to each other only loosely if at all. Happily, in Vermont Prof. Shimony has actually found Machs’ General Store (Fig. 1), and characterizes it thus: “what we have is on the shelves; what you don’t see don’t exist; we give no credit.”

The dragging effects are probably the most prominent items in Mach’s store, and Pfister gives their interesting history. One could supplement this with the history of other Machian effects, because one use of a good principle is to make simple and quick qualitative or semi-quantitative suggestions for a variety of different physical situations. The most fruitful and imaginative use of Mach’s principle in this fashion occurred in the 50’s and 60’s, largely inspired by Dicke. Some of the resulting ideas have appeared in the literature (for example, Dicke 1961, 1963, Peebles and Dicke 1962), but they found their

most creative expression in Dicke's informal "brainstorming" sessions for the Princeton relativists. Here all sorts of ways that the universe might influence local physics were explored, and the magnitudes of possible experimental observations were determined by liberal use of Mach's principle.

2. Dragging of inertial frames

Of all the predictions that follow from, or have been read out of, Mach's principle, the dragging of inertial frames by rotating bodies is certainly the most definite and least controversial. If one measures this dragging by the Coriolis forces — that is, one calculates in the first order of the angular velocity — then the answer of General Relativity is unambiguous, and can be derived simply and elegantly (Pfister 1993). This answer can be interpreted to agree with the Machian expectations, at least if one is willing to refine them suitably and to some extent after the fact. Mach himself, in his famous passage (Mach 1883), simply informs us that the dragging might be measured by centrifugal forces, but it is reasonable to include Coriolis forces (and other effects!) with respect to Mach's principle. After all, no one is competent to say how Mach's principle would have turned out if the background of its author increased in scope and style till it ultimately encompassed several modern alternative theories of gravity.

It is the particular merit of Pfister and his students that they did a detailed study of the Machian centrifugal force terms. By the "correct" centrifugal force Pfister means the one that corresponds to the spacetime region being *flat*. As he remarks, to have a centrifugally correct region, flat to second order in the angular velocity, necessarily demands a rather special situation: nonlinearity will generally cause second-order forces other than pure centrifugal ones. Thus he finds that the correct forces inside a shell require that the shell have special properties, such as a prolate shape, differential rotation, etc.

But given these special properties one *can* induce the "correct" inertial forces for the case of rotation; in fact, Pfister conjectures that the effects of *arbitrary* acceleration can be induced in a laboratory at rest with respect to the fixed stars by suitably moving exterior masses. This is a very interesting conjecture, not least because it forces us to think about the proper definition of global measurements, such as acceleration with respect to the fixed stars, and about other global Machian quantities. For example, naively one might reject the conjecture because the inertial forces are "coordinate effects," whereas the effect of exterior masses is presumably a curvature effect; but this would neglect the global nature of the acceleration that is at issue here.

In the general situation, when the exterior masses are not moving "suitably," Pfister shows that it is difficult to distinguish between Machian dragging and gravitational radiation. If we say that it is in fact impossible, we come very close to the type of minimalistic view of Mach's principle advocated for example in the Mach-Einstein-Wheeler formulation (Isenberg 1993). But just because the Machian dragging is a (global) "coordinate effect," whereas gravitational radiation is a (local) "curvature effect" we can equally well view the difficulty as a challenge, to define the Machian dragging in a general and useful way so that it *can* be distinguished from gravitational radiation.

3. Alternative Theories

This is not the place to review the many ways in which Mach's principle has suggested alternative theories. In the case of inertial dragging, Mach of course makes no explicit prediction that might confirm or contradict the general relativity result; but in the limit when the rotating mass is the whole universe, the usual expectation is that of "complete dragging." It may be that in this limit the small-rotation results will be exact (because complete dragging presumably means that the rotation with respect to the inertial frame is vanishingly small). As a first step toward a proof of such conjectures nothing would be more helpful than a clear, general definition of "complete dragging."

If this cosmological Machian expectation is not satisfied, it may be the fault of the cosmological model. The task would then be to find a more restricted class of models that are Machian, as advocated in a number of formulations (such as the Einstein-Wheeler-Mach version). Or it may be the fault of the theory, in which case one would look for a more Machian theory of gravity; this is one of the claims of the Brans-Dicke theory. Calculations of dragging effects in alternative theories are appropriate not only to determine this "degree of Machismo," but also because these effects will eventually become measurable experimentally, and it is interesting to know how accurately one will have to measure in order to distinguish alternative theories on the basis of dragging. This question has been answered (to lowest order) in the *PPN* formalism, but beyond that the literature appears limited.

Two alternative theories are particularly interesting in this context, Brans-Dicke theory (because it takes Mach's principle as a motivational basis) and low-energy string theory (because string theory is supposed to be more fundamental than general relativity). Both of these incorporate a scalar field ("dilaton") as part of the gravitational force. That the scalar part of gravity should not contribute to dragging is suggested by symmetry considerations: to lowest order, rotational perturbations will affect only the g_{0i} components of the metric, and *not* the one component of the scalar field. Therefore in Brans-Dicke theory, for example, the dragging is reduced compared to general relativity (Brill 1962), because the scalar field contributes positively to the gravitational force but "does not drag." One can also show that this theory is, in a sense, *more* Machian than general relativity: it leads to the limit of complete dragging in the limit when the rotating shell is the only matter in the universe, independent of the shell's mass: the gravitational "constant" will automatically adjust so that the shell is at its Schwarzschild radius.

Dragging effects appear not to have been discussed in the literature on low-energy string theory, which has no natural place for phenomenologically described matter. The long-range fields in this theory are the gravitational, scalar (dilaton), electromagnetic, and axion fields. However, one does have solutions to this theory for rotating black holes (Sen 1992). For uncharged black holes these show *no* difference from general relativity, and appear to suggest that the dragging is not reduced as in Brans-Dicke theory. This may be surprising if one expects the rotating black hole solutions to reproduce, at least to lowest order, the exterior dragging effects of other rotating configurations; but the reason is easily explained.

The black hole solutions of low-energy string theory are the simplest solutions to the *sourceless* equations corresponding to black holes. In these equations the electromagnetic

field generates the scalar field; if the former vanishes (zero charge), the scalar field is constant. In the Brans-Dicke theory, on the other hand, one has in mind matter sources, and the scalar field is generated by the trace of the matter's stress-energy tensor. Thus the difference already occurs in the description of an unperturbed mass center: Brans-Dicke theory attributes a part of the gravitational force from matter to the scalar field, whereas low-energy string theory considers the scalar field to be constant for uncharged black holes. In fact, collapse to a black hole in Brans-Dicke theory also leads to black holes with a constant scalar field, which are the same as those of general relativity (Hawking 1972). Thus there is no reason to suppose that the role of the scalar field in dragging effects is different in these theories. (Whether there are also other types of black holes in Brans-Dicke theory is the subject of current research, Campanelli and Lousto 1993).

Footnote

*Chapter 6, “Der Blick ins Große — zum Scheitern verurteilt?”

Figure caption

Fig. 1: Machs' General Store, Pawlet, Vermont

References

- Brans, Carl H. and Dicke, Robert H. “Mach's Principle and a Relativistic Theory of Gravitation” *Physical Review* **124** 925-935
- Campanelli, M. and Lousto, C. O. “Are Black Holes in Brans-Dicke Theory precisely the same as in General Relativity?” *preprint* Universität Konstanz; gr-qc/9301013
- Dicke, Robert H. 1961 “Experimental Tests of Mach's Principle” *Physical Review Letters* **7** 359-360
- Dicke, Robert H. 1963 “Cosmology, Mach's Principle and Relativity” *American Journal of Physics* **31** 500-509
- Dicke, Robert H. 1962 “The Earth and Cosmology” *Science* **138** 653-664
- Fahr, Hans Jörg 1993 *Der Urknall kommt zum Fall* Frankh-Kosmos
- Hawking, K. Stephen 1972 “Black Holes in the Brans-Dicke Theory of Gravitation” *Communications in Mathematical Physics* **25** 167-171
- Isenberg, James 1993 “On the Wheeler-Einstein-Mach Spacetimes” this volume
- Peebles, P. James and Dicke, Robert H. 1962 “Significance of Spatial Isotropy” *Physical Review* **127** 629-631
- Pfister, Herbert 1993 “Dragging effects near rotating bodies and in cosmological models” previous paper, this volume
- Sen, A. 1992 “Rotating Charged Black Hole Solutions in Heterotic String Theory” *Physical Review Letters* **69** 1006
- Shimony, Abner 1992 *private communication* APS Spring Meeting, Washington, DC

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9310033v1>